




Bound-free and bound-bound spectroscopy of cold trapped molecular ions



Roland Wester
**71st International Symposium on
Molecular Spectroscopy**
Urbana-Champaign
June 21, 2016

Institut für Ionenphysik und Angewandte Physik
Universität Innsbruck
Austria







Outline

- ❖ Spectroscopy in multipole rf ion traps
- ❖ Vibrational overtone transitions
- ❖ Bound-free photodetachment spectroscopy
- ❖ Terahertz rotational spectroscopy
- ❖ Electronic spectra of the Atkins' snowball

Multipole rf ion traps

The 22-pole ion trap


Dieter Gerlich, Physica Scripta (1995)



- Large almost field free volume
 - Variable temperature 8-300 K
 - Buffer gas cooling of internal and translational motion
 - Neutral reactants @ well defined density
- RW, J. Phys. B (2009)



22-pole trap setup based on rf mass spectrometers



“Classical” 22-pole trap setup by D. Gerlich et al.




Spectroscopy by laser-induced reactions






Spectroscopy of H_3^+ by laser-induced reactions

Vibrational levels of H_3^+




breathing mode



Infrared
inactive

degenerate
bending modes



Infrared active



Jochen Mikosch, Holger Kreckel,
Radek Plasil, Juraj Glosik,
Dieter Gerlich, Andreas Wolf, Dirk
Schwalm, RW, J. Chem. Phys. (2004)



Spectroscopy of H_3^+ by laser-induced reactions






The lowest rotational states

$$E_{\text{rot}} = \hbar^2 \frac{J(J+1)}{2 I_{\perp}} + \frac{K^2 \hbar^2}{2} \left(\frac{1}{I_{\parallel}} - \frac{1}{I_{\perp}} \right)$$






Spectroscopy of H_3^+ by laser-induced reactions



Count ArH^+
products!




Spectroscopy of H_3^+ by laser-induced reactions






Spectroscopy of H_3^+ by laser-induced reactions



Spectroscopy of H_3^+ by laser-induced reactions






Outline

- ❖ Spectroscopy in multipole rf ion traps
- ❖ Vibrational overtone transitions
- ❖ **Bound-free photodetachment spectroscopy**
- ❖ **Terahertz rotational spectroscopy**
- ❖ Electronic spectra of the Atkins' snowball




Rotational transitions in cold OH⁻



Couplings between rotational states:

- Collisions with helium buffer gas
- Terahertz rotational transitions


Photodetachment as probe of J



$$\Delta E = 37.5\text{cm}^{-1} \text{ or } 54 \text{ K}$$





The Innsbruck 22-pole trap setup






Photodetachment spectroscopy of cold OH⁻






Rotational transitions in cold OH⁻



Couplings between rotational states:

- Collisions with helium buffer gas
- Terahertz rotational transitions


Photodetachment as probe of J



$$\Delta E = 37.5\text{cm}^{-1} \text{ or } 54 \text{ K}$$

Rotational transitions in cold OH⁻

Photodetachment with *strong* laser



$$\frac{d}{dt}N_0 = -k_{01} \cdot N_0 + k_{10} \cdot N_1$$

$$\frac{d}{dt}N_1 = +k_{01} \cdot N_0 - [k_{10} + k_{PD}] \cdot N_1$$

→ Laser depletion of excited state yields inelastic collision rate



k_{10}, k_{01} : helium inelastic collision rates



OH⁻ rotational quenching: J=1→0

Changing the rotational state population

Ion collision temperature: 22 K




Faster detachment rate → stronger depletion of J=1

Fit: model with inelastic collision rate k_{01} as only free parameter

OH⁻ rotational quenching: J=1→0

Density dependent inelastic rates

Ion collision temperature: 22 K



Slope: absolute inelastic rate coefficient


OH⁻:
 $k_{J=1 \rightarrow 0} = (1.8 \pm 0.4_{\text{stat}} \pm 0.2_{\text{syst}}) \times 10^{-11} \text{ cm}^3/\text{s}$

OD⁻:
 $k_{J=1 \rightarrow 0} = (2.8 \pm 1.1_{\text{stat}} \pm 0.4_{\text{syst}}) \times 10^{-11} \text{ cm}^3/\text{s}$



OH⁻ / OD⁻ rotational quenching: J=1→0

Quantum scattering calculations



Time-independent quantum coupled-channel calculations

- Full Born-Oppenheimer surface, identical for OH⁻ and OD⁻ MP4/aug-cc-pVQZ
- Partial waves up to 40h

→ Substantial differences for OH⁻ and OD⁻


Fabio Carelli & Franco Gianturco

D. Hauser *et al.*, Nature Physics (2015)



OH⁻ rotational quenching: J=1→0



Thermal inelastic rate coefficients






Direct OH⁻ rotational excitation

Terahertz spectroscopy with difference frequency mixing




Antenna output power





Direct OH⁻ rotational excitation


Bringing the THz antenna close to the ion trap





Direct OH⁻ rotational excitation

Terahertz spectroscopy via photodetachment




Important:
Low rates for collisional
excitation/deexcitation



Direct OH⁻ rotational excitation

Terahertz spectroscopy via photodetachment







Outline

- ❖ Spectroscopy in multipole rf ion traps
- ❖ Vibrational overtone transitions
- ❖ Bound-free photodetachment spectroscopy
- ❖ Terahertz rotational spectroscopy
- ❖ **Electronic spectra of the Atkins' snowball**



Electronic spectra of an Atkins' snowball

(an experiment
without an ion trap)



H. Linnartz, X. Tielens, J. Cami, A. Y. Wang, M. Alcamí,
F. Martín, M. Beyer, RW, A. Lindinger, *et al.*
P. Scheier

THE PHYSICAL REVIEW

A journal of experimental and theoretical physics established by E. L. Nichols in 1893

SECOND SERIES, VOL. 116, No. 6

DECEMBER 15, 1959

Ions in Liquid Helium*

K. R. ATKINS


Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania

(Received July 16, 1959; revised manuscript received September 30, 1959)



Electronic spectra of an Atkins' snowball


$C_{60}^+(He)_n$ formation in He droplets



Summary

Spectroscopy in ion traps by laser-induced reactions

:: Vibrational spectroscopy of H_3^+ overtones



Rotational spectroscopy of trapped ions



:: Near-threshold detachment for $\text{OH}^- \rightarrow$ rotational populations

:: Cold state-to-state inelastic rate coefficients

:: Rotational excitation by direct THz absorption

Electronic spectra in He droplets

:: Line shifts of $\text{C}_{60}^+(\text{He})_n$ Atkins snowball reveal phase transitions




Outlook

:: THz spectroscopy of polyatomic ions

:: Dipole-bound states in carbon chain ions

:: Inelastic collision rates for open-shell ions





Thank you

Current group

Olga Lakhmanskaya (Ph.D. st.)
Eric Endres (Ph.D. st.)
Steffen Spieler (Ph.D. st.)
 Ibrokhim Iskandarov (Ph.D. st.)
Katharina Geistlinger (Ph.D. st.)
 Tim Michaelsen (Ph.D. st.)
 Malcom Simpson (Ph.D. st.)
 Björn Bastian (Ph.D. st.)
 Atilay Ayasli (Master st.)
Lena Remmers (Master st.)
Moritz Fischer (Master st.)
Fabio Carelli (Postdoc)
 Mario Hernandez (Postdoc)
 Jennifer Meyer (Postdoc)
 Martin Ruetz (Electronics Eng.)
Franco A. Gianturco (Sen. Prof.)
 Ingeborg Rauter (Secr.)



www.uibk.ac.at/ionen-angewandte-physik/molsyst

Former group members

Eduardo Carrascosa
 Gerhard Egger
 Martin Stei
Daniel Hauser
Seunghyun Lee
 Martin Bawart
 Martin Kainz
Sunil Kumar
 Alexander Doerfler
Thorsten Best
 Josef Wildauer
 Aditya Kelkar
 Fabian Hochheimer
 Johannes v. Vangerow
Rico Otto
 Johannes Deiglmayr
 Sebastian Trippel
 Petr Hlavenka
 Jochen Mikosch

Collaborations

Matthias Weidemüller (U Heidelberg)
 Bill Hase (Texas Tech Univ.)
 Stepan Roucka & Juraj Glosik (U Prague)
 Wolf Geppert (U Stockholm)
 Tom Millar (U Belfast)
 Olivier Dulieu (CNRS Orsay)
 Dave Parker (U Nijmegen)
 Paul Scheier (U Innsbruck)

